


**COMPLIANCE
RE-IMAGINED**

Preventing Clinical Errors in Practice


Scott Munsterman, DC, FICC, CPCO



Scott Munsterman, DC, FICC, CPCO Brief Bio

Dr. Scott Munsterman is an acknowledged expert on the transforming model of health care delivery and compliance with a commitment to the promotion and advancement of the chiropractic profession. Dr. Munsterman is founder and CEO of Best Practices Academy, a clinical improvement organization providing focused leadership to bring practices into compliance with regulatory standards, equip them to improve clinical outcomes, and integrate into the transformed care delivery system. Dr. Scott works with ClinicArmor and BPA's EHR partnered with iPatientCare.

Dr. Munsterman is a graduate of Northwestern Health Sciences University, where he has served as Vice-Chair of the Board of Trustees and on the President's Cabinet as Chief of Care Delivery. He was awarded Chiropractor of the Year in South Dakota and the Fellow of the International College of Chiropractors (FICC). He is a professional compliance officer. Dr. Munsterman served two terms as Mayor of the City of Brookings and three consecutive terms in the South Dakota House of Representatives, where he chaired the House Health and Human Services Committee and also chaired the Legislative Planning Committee. He is author of the books "A Vision for South Dakota", "Care Delivery and Chiropractic: An Opportunity Waiting", and "Unfinished Business". However, he states his greatest accomplishment has been his five daughters and six grandchildren - with more success to come.




Disclaimer

The topics taught here are for the sole purpose of the chiropractic profession, any transference to other healthcare disciplines are at the risk of the individual's discretion. The presenter is an investor in the Best Practices Academy and ChiroArmor/ClinicArmor. The Best Practices Academy and ChiroArmor/ClinicArmor denies responsibility or liability for any erroneous opinions, analysis, and coding misunderstandings on behalf of individuals undergoing this course.

This presentation was current at the time it was published or uploaded onto the web. Medicare policy changes frequently so links to the source documents have been provided within the document for your reference. We have based the majority of this program on the guidelines set forth by the OSHA, OCR, HHS, CMS, NCCQA, URAC, AAAHC, AHRQ, and other agencies involved in health care standards and research dissemination, as it relates to the chiropractic profession. We encourage readers to review the specific statutes, regulations, and other interpretive materials for a full and accurate statement of their contents.

No legal advice is given in this program, and we encourage you to refer any such questions to your healthcare attorney.




1-3

<http://www.bestpracticesacademy.com>




What is Patient Safety?

First do no harm.




Office of Inspector General
Report in Brief
May 2022, OIG-06-18-10000

25% of Medicare patients experience patient harm during their hospital stay.

Patient harm includes adverse events and temporary harm events.

- 12 percent** of patients experienced adverse events, which are events that led to longer hospital stays, **permanent harm**, life-saving intervention, or death.
- 13 percent** of patients experienced **temporary harm** events, which required intervention but did not cause lasting harm, prolong hospital stays, or require life-sustaining measures.



Physician-reviewers determined that 43 percent of the harm events could have been prevented if patients had been provided better care.



Copyright 2011, Cleveland. All Rights Reserved.

56% of harm events were not preventable and occurred even though providers followed proper procedures...



Copyright 2011, Cleveland. All Rights Reserved.

Terminology



Copyright 2011, Cleveland. All Rights Reserved.

Patient safety: the avoidance, prevention and amelioration of adverse outcomes or injuries stemming from the process of health care.

National Patient Safety Foundation. Agenda for research and development in patient safety.
<http://www.ih.org/Topics/PatientSafety/Pages/default.aspx>



Copyright 2011, Cleveland. All Rights Reserved.

What is an Adverse Outcome or Event?

An **unexpected and undesired incident** directly associated with the care or services provided to the patient; an incident that occurs during the process of providing health care and **results in patient injury or death; or an adverse outcome for a patient, including an injury or complication.**



Copyright 2011, Cleveland. All Rights Reserved.

Preventing Clinical Errors

A Clinical Error is an **act of omission or commission in planning or execution** that contributes or could contribute to an unintended result.

Defining medical error. Ethan D. Grober, John M.A. Bohnen Can J Surg. 2005 Feb; 48(1): 39-44. PMID: PMC3211566
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3211566/>



Copyright 2011, Cleveland. All Rights Reserved.

Preventable Harm

The Institute for Healthcare Improvement defines preventable medical harm as **“unintended physical injury resulting from or contributed to by medical care (including the absence of indicated medical treatment)”**, that requires additional monitoring, treatment or hospitalization, or that results in death.”

Recent studies (as of 2017) of medical errors have estimated errors may account for as many as **251,000 deaths annually** in the United States (U.S.), making medical errors the **third leading cause of death**

Anderson JG, Abrahamson K. Your Health Care May Kill You: Medical Errors. Stud Health Technol Inform. 2017;234:13-17. PMID: 28186008.



Copyright 2021. All Rights Reserved.


Causes of Errors

Adverse Events vs Near Misses
Human vs System
Commission vs Omission



Copyright 2021. All Rights Reserved.

NCC MERP Index for Categorizing Medication Errors



© 2021 National Coordinating Council for Medication Error Reporting and Prevention. All Rights Reserved.

* Permission is hereby granted to reproduce information contained herein provided that each reproduction shall not modify the text and shall include the copyright notice appearing on the pages from which it was copied.



Copyright 2021. All Rights Reserved.

Most errors are the result of various causes and predisposing conditions.

In other words, there are a variety of factors involved that can lead to or cause a clinical error or adverse event – or a near miss.



Copyright 2021. All Rights Reserved.

Swiss Cheese Model


Reason J. Human error: Models and management. BMJ 2000; 320:768-70



Copyright 2021. All Rights Reserved.


Holes in the Defense Layers

A bad outcome occurs only when the holes in many defense layers momentarily line up to permit a trajectory of an accident opportunity—bringing hazards into damaging contact with patients.



Copyright 2021. All Rights Reserved.

It just takes one thing to block the incident...



Copyright © 2011, Elsevier. All Rights Reserved.

Reasons for Holes in the Defense Layers

Active Failures are the unsafe acts committed by people who are in direct contact with the patient or system. They take a variety of forms: slips, lapses, fumbles, mistakes, and procedural violations.

Latent Conditions have two kinds of adverse effects:

- they can translate into error provoking conditions within the workplace (i.e. time pressure, understaffing, inadequate equipment, fatigue, inexperience) and
- they can create long-lasting holes or weaknesses in the defenses (i.e. lack of training for staff, improper therapeutic or billing practices, lack of compliance policy).



Copyright © 2011, Elsevier. All Rights Reserved.

Types of Clinical Errors

- Diagnostic
- Treatment
- Preventive
- Other

National Academies of Sciences, Engineering, and Medicine. 2015. *Improving diagnosis in health care*. Washington, DC: The National Academies Press.



Copyright © 2011, Elsevier. All Rights Reserved.

Diagnostic Error

"the failure to establish an accurate and timely explanation of the patient's health problem(s) or communicate that explanation to the patient."

National Academies of Sciences, Engineering, and Medicine. 2015. *Improving diagnosis in health care*. Washington, DC: The National Academies Press.



Copyright © 2011, Elsevier. All Rights Reserved.

57% of all diagnostic failures occur in ambulatory care settings.




Copyright © 2011, Elsevier. All Rights Reserved.

Diagnostic Process: 7 Stages

1. Access and presentation
2. History taking/collection
3. Physical exam
4. Testing
5. Assessment (differential diagnosis)
6. Care planning/referral
7. Follow-up/Outcome Assessment



Copyright © 2011, Elsevier. All Rights Reserved.



Methods for Detecting Failure Across the Diagnostic Process

Work System/Environmental Factors
Common Points of Failure

Work System/Environmental Factors which may produce failure in the Diagnostic Process

1. Information gathering and integration/communication (amount, accuracy, completeness, appropriateness)
2. Technology (EHR is the right fit, full adoption into workflows)
3. Organization and roles of providers and staff
4. Physical space and layout

Common Points of Failure in the Diagnostic Process

1. Patient delay in seeking care
2. Failure to gather enough information
3. Failure to integrate the information in medical decision-making process
4. Failure to establish an accurate diagnosis
5. Failure to communicate an explanation/diagnosis to the patient

Standardization

Monitor and adjust to changes, make improvements, build-in check systems.

Key Areas to Improve Diagnostic Performance

- Clinical reasoning
- Teamwork
- Communication with patients, their families, and other health care professionals
- Appropriate use of diagnostic tests and the application of these results on subsequent decision making
- Use of health IT

Clinical Reasoning



Copyright 2021, Drexel University. All Rights Reserved.

Clinical Reasoning:

Understand Heuristics and Biases that Influence Decision Making

- **Anchoring:** tendency to "lock" onto features of the initial presentation and failure to adjust this initial impression in light of new information.
- **Affective Bias:** letting our emotions, feelings, and biases affect our judgement.
- **Availability Bias:** tendency to more easily recall certain things that have been seen recently or that are common or impressed upon us.
- **Context Errors:** instances where we misinterpret the situation, leading to an erroneous conclusion.
- **Search Satisficing:** premature closure resulting from accepting the first answer that comes along that explains the facts without considering whether there might be a different or better solution.



Copyright 2021, Drexel University. All Rights Reserved.

Dual Process Theory



Copyright 2021, Drexel University. All Rights Reserved.

Dual Process Theory and Diagnosis

- **System 1:** When a patient presents, the initial data include **typical symptoms and signs of disease** which are recognized. System 1 processing is fast.
- **System 2:** When the symptoms and signs are **atypical and do not become apparent to align with a specific disease pattern**. Repetition of data to System 2 may eventually be recognized as a new pattern and then processed through System 1. System 2 processing is slow, analytical decision making and can override System 1 processing.

Expert clinicians possess better developed mental models of diseases, which support more reliable pattern matching (System 1). Novice clinicians are more likely to rely on analytical reasoning throughout the diagnostic process as compared to experienced clinicians.



Copyright 2021, Drexel University. All Rights Reserved.

What about our staff and the role they can play in preventing diagnostic errors?



Copyright 2021, Drexel University. All Rights Reserved.

Teamwork

- Gathering information is essential
- It takes a team to monitor and screen patients on an ongoing basis...
- Clarify the roles of each team member within the practice and their responsibilities
- Assess their competence within their roles.
- Professionalism
- Clinical Conscientiousness and Situational Awareness



Copyright 2021, Drexel University. All Rights Reserved.

Communication

Providers and staff collaborating with their experience with the patient and communication efforts.



Copyright © 2021 Oakhurst. All Rights Reserved.

Use of Diagnostic Tests

What is clinically indicated based on standard of care?



Copyright © 2021 Oakhurst. All Rights Reserved.

Use of Health IT

Has the practice fully adopted an EHR which meets the compliance and patient documentation, clinical decision making needs of the type of patient care delivered?



Copyright © 2021 Oakhurst. All Rights Reserved.

Treatment

Is/was the procedure being performed correctly?
Is equipment (therapeutic modalities, etc.) functioning properly?



Copyright © 2021 Oakhurst. All Rights Reserved.

Preventive: Ongoing Assessment

Ask yourself these questions...



Copyright © 2021 Oakhurst. All Rights Reserved.

Patient Assessment:

Key Triggers to Identify Patients At-Risk for Adverse Events

- Is the patient taking medication?
- Has the patient recently had surgery or another medical intervention?
- Does the patient show any signs of a change in mental status?
- Does the patient demonstrate any unsteadiness in walking, standing, or movement – such as sitting to standing?
- Has the patient had any falls within the past 12 months?
- Is the patient allergic or have a sensitivity to anything?
- Does the patient have multiple comorbidities and/or a poor health status?



Copyright © 2021 Oakhurst. All Rights Reserved.

Patient Assessment:

Key Triggers to Identify Patients At-Risk for Adverse Events

- What has changed since the last visit for the patient (monitoring response)?
- Any new information to integrate into current working diagnosis (contraindications, red flags ruled-out)?
- Clinical reasoning and treatment response reflect certainty in diagnosis?
- Communication with patient reflects engagement?



Copyright © 2021 Chiropractor. All Rights Reserved.

What are the defense layers in the practice?

1. Emergency identification/response procedures are in place.
2. Performing vital signs.
3. Properly diagnosis a patient's condition.
4. Identifying contraindications for care and red flags.
5. Perform manipulation procedure properly.
6. Safely apply therapeutic procedures/activities on each visit.
7. Close oversight/response of patient monitoring during care.
8. Close oversight of visitors/children during patient's visit.
9. Awareness of external activities within and outside of the facility.
10. Doctor/Staff rested and devote 100% present time consciousness.



Copyright © 2021 Chiropractor. All Rights Reserved.

Other

- Equipment failure
- Communication failure
- Failure in systems process, workflows, etc.



Copyright © 2021 Chiropractor. All Rights Reserved.

If an incident does occur...

- The incident should not be kept secret. All incidents need to be documented and discussed with your professional liability insurer – and then with other providers and staff.
- The doctor should talk to the patient (if recommended by your professional liability insurer):
 - Discuss what has been learned
 - Provide an honest expression of regret or apology
 - Can often decrease the risk of legal action



Copyright © 2021 Chiropractor. All Rights Reserved.

How can we address patient safety in our practice?



Copyright © 2021 Chiropractor. All Rights Reserved.

The “Walk”

Welcome
Ask
Listen
Knowledge



Copyright © 2021 Chiropractor. All Rights Reserved.

Screening Patients:

Why are you here today?
Has there been a change in how you are feeling since your last visit?
Have you seen anyone else about your health?
Do you have questions about...
Are you worried about your health?

Situational Awareness:

No change or worsening
Observation of patient's mental status, behaviors, or characteristics

Has there been a "Significant Event"?



Copyright © 2021 Cleveland. All Rights Reserved.

Patient Safety Questionnaire

Patient Safety Questionnaire
Please answer the following questions as it pertains to your visit today - we will use this information as we strive to help you!

- | Yes | No | |
|--------------------------|--------------------------|---|
| <input type="checkbox"/> | <input type="checkbox"/> | I have felt unsteady and off balance when I walk, stand or move. |
| <input type="checkbox"/> | <input type="checkbox"/> | I have recently had a hospitalization or surgery since my last visit. |
| <input type="checkbox"/> | <input type="checkbox"/> | There has been a change in my condition since my last visit. |
| <input type="checkbox"/> | <input type="checkbox"/> | Something happened to me since my last visit (i.e., accident, fall, etc.) |
| <input type="checkbox"/> | <input type="checkbox"/> | I have had a recent change in my medication or have a new allergy. |
| <input type="checkbox"/> | <input type="checkbox"/> | I have fallen within the last 6 months. |
| <input type="checkbox"/> | <input type="checkbox"/> | My mental processing or thinking doesn't seem very clear or is different to me. |
| <input type="checkbox"/> | <input type="checkbox"/> | I need to talk to the doctor about my progress. |
| <input type="checkbox"/> | <input type="checkbox"/> | I am concerned about the treatment I am getting. |
| <input type="checkbox"/> | <input type="checkbox"/> | I have questions I need to ask the doctor today. |



Copyright © 2021 Cleveland. All Rights Reserved.

Does the patient's clinical presentation require urgent need for evaluation and/or care?

The doctor must be informed of any new information about the patient that has been related to staff.



Copyright © 2021 Cleveland. All Rights Reserved.

Stay Connected to Established Patients who are under a treatment plan.

Following the treatment plan, evidence-informed care guidelines, and the patient's response to care...



Copyright © 2021 Cleveland. All Rights Reserved.

Screening Patients

Monitor changes since the last visit
No change or worsening
Observation of patient's behaviors and characteristics



Copyright © 2021 Cleveland. All Rights Reserved.

Has there been a "Significant Event"?



Copyright © 2021 Cleveland. All Rights Reserved.

Does the patient's clinical presentation require urgent need for evaluation and/or care?

The doctor must be informed of any new information about the patient that has been related to staff.



Copyright © 2021 Osha/HRSA. All Rights Reserved.

Stay Connected to Established Patients who are under a treatment plan.

Following the treatment plan, evidence-informed care guidelines, and the patient's response to care...



Copyright © 2021 Osha/HRSA. All Rights Reserved.

Patient Safety is First and Foremost

There should be an ongoing discussion regarding strategy towards preventing clinical errors and enhancing patient safety.



Copyright © 2021 Osha/HRSA. All Rights Reserved.

Most Common Patient Safety Issues

- Falls
- Equipment malfunction
- Infection prevention procedures
- Faulty patient perception of an incident occurring stemming from lack of communicating to the patient what to expect from treatment
- Underlying medical emergency/red flag (i.e., cardiovascular, cerebrovascular, fracture, infection, cancer)



Copyright © 2021 Osha/HRSA. All Rights Reserved.

Recognizing Patient Safety Incidents

- Patient complains of pain after treatment
- Modality malfunctioning or not being applied properly
- Patient nearly falling
- Patient safety incidents range from "No Harm" to "Unnecessary Harm"



Copyright © 2021 Osha/HRSA. All Rights Reserved.

Underlying Causes


- Patient OTC drug use and interactions increase risk of falls
- Provider/therapist fatigue and stress can lead to miscommunications
- Short staffing and increased workload



Copyright © 2021 Osha/HRSA. All Rights Reserved.

Follow Safe Practice Procedures

You and your staff must be the **"one thing"**...



Copyright © 2021 ChiroForum. All Rights Reserved.

Scope of Safety Issues

In Chiropractic...



Copyright © 2021 ChiroForum. All Rights Reserved.

Professional Boundaries in Clinical Practice

Patient Relationships



Copyright © 2021 ChiroForum. All Rights Reserved.

Professional boundaries are limits which protect a worker's professional power and their patient's vulnerability. Successful and ethical working relationships are based on a clear understanding of what the workers' role is – and just as importantly – what their role isn't.

Definition of Professional Boundary

<https://mcarthur.com.au/media/1429/understanding-professional-boundaries.pdf>



Copyright © 2021 ChiroForum. All Rights Reserved.

When does Clinical Integrity become compromised?

THE ROLE	THE LINE	THE IMPACT
Professional Service		
Social Interaction		
Mutual Friendship	BOUNDARY CROSSING	CONSCIOUSNESS OF BOUNDARY VIOLATION PENDING
Close Friendship	BOUNDARY VIOLATION	PERSONAL GAIN
Family		EMOTIONAL DEPENDENCY
Intimacy		VIEWED AS EXPLOITATION IF PROFESSIONAL ROLE IS CONTINUED



ChiroForum

What is our role as a health care professional?

- Perform clinical duties and provide care to a patient
- Protect the patient from harm
- Meet reasonable expectations of the patient
 - Respect and dignity
 - Provide competent care
 - Practice ethically
 - Uphold confidentiality
 - Comply with all laws regulating your practice and behaviors
- Honesty in all patient interactions
- Equitable and fair treatment of all patients regardless of their race, religion, socioeconomic status, etc.



Copyright © 2021 ChiroForum. All Rights Reserved.

Has your allegiance shifted away from your focus in your professional role to a more personal role whereby you are seeking and benefiting personally from the relationship?



Copyright © 2021, Elsevier Inc. All Rights Reserved.

Commonly Misdiagnosed Conditions

The "Big Three": misdiagnosed cancers (37.8%), vascular events, like stroke and heart attack (22.8%), and infections (13.5%).

Cancers

Lung, breast, colorectal, prostate, and skin cancers

Vascular events

Stroke, heart attack, venous thromboembolism (blood clots in the legs and lungs), aortic aneurysm and rupture (dissection), arterial thromboembolism (a blockage of the blood supply to internal organs)

Infections

Sepsis, meningitis, encephalitis, spinal infection, pneumonia, and endocarditis (a heart infection)

Newman-Tokoh, D. L., Schaffer, A. C., Yu, M., C. Nasser, N., Sabar, Tattari, A. S., Clemons, G. D., Wang, Z., Zhu, Y., Faria, M., & Saggi, D. (2019). Serious misdiagnosis-related harms in inpatient claims: The "Big Three" - vascular events, infections, and cancers. *Diagnosics*, 4(2), 227-240. doi: <https://doi.org/10.1016/j.diag.2019.05.004>



Copyright © 2021, Elsevier Inc. All Rights Reserved.

What are the various factors that may set us up for risk of a clinical error in practice?



Copyright © 2021, Elsevier Inc. All Rights Reserved.

Evaluation Process



Copyright © 2021, Elsevier Inc. All Rights Reserved.

Intake Forms

Accurate information from the patient is imperative to gather for your doctor's clinical decision making.



Copyright © 2021, Elsevier Inc. All Rights Reserved.

Reasons for Diagnostic Error

Patients are interrupted when telling their illness story to providers within 11-18 seconds.^{2,3}



This breakdown in collaborative communication may result in assumptions and premature closure.⁴



These breakdowns lead to diagnostic error in about 1 of every 20 patients you see.⁵



How Does 60 Seconds To Improve Diagnostic Safety Work?

Ask

Listen

Act

AHRQ Agency for Healthcare Research and Quality Advancing Excellence in Health Care | www.ahrq.gov

Tips for Effective Listening

- Listen for meaning.
- Pay attention to body language.
- Cultivate empathy.
- Avoid making judgments.
- Look into others' eyes when they're speaking.
- Pay attention to the feelings associated with the words.

AHRQ Agency for Healthcare Research and Quality Advancing Excellence in Health Care | www.ahrq.gov

How can we improve diagnostic safety?

What can patients do?

- ✓ Tell their story fully and completely and clearly
- ✓ Provide accurate information about their symptoms
- ✓ Speak up if they feel they have not been heard
- ✓ Ask questions to clarify the information shared
- ✓ Use a checklist of tests, symptoms, concerns, or physicians consulted

What can clinicians do?

- ✓ Listen to patients
- ✓ Support patients in effectively sharing their symptoms.
- ✓ Ask patients what they think is going on
- ✓ Conduct a thorough history and physical examination
- ✓ Set a visit agenda
- ✓ Know patients and their history, and read prior notes
- ✓ Integrate "pre-work" for patients (e.g. symptoms; history of present illness; labs)

AHRQ Agency for Healthcare Research and Quality Advancing Excellence in Health Care | www.ahrq.gov

HOW IT WILL WORK IN YOUR PRACTICE?

Initial Visit Screening

Initial Visit Patient Safety Checklist

Use this checklist to gather essential information regarding the initial or the admission of a patient to determine if the patient is at risk for an adverse event and there is a need for the monitoring of the patient safety activities.

PATIENT INTAKE AND INFORMATION GATHERING	
<input type="checkbox"/> Intake forms completed and signed	<input type="checkbox"/> Patient is showing understanding in responses
<input type="checkbox"/> Patient's mental status is not normal	<input type="checkbox"/> Patient has fallen within the last 8 months
<input type="checkbox"/> Patient recently had a hospitalization or surgery	<input type="checkbox"/> Patient is on medication or has allergies

PRE-EVAL EHR DOCUMENTATION	
<input type="checkbox"/> Demographic completed	<input type="checkbox"/> Therapies documented and communicated
<input type="checkbox"/> Referring Physician identified and documented	<input type="checkbox"/> Patient's Chief Complaint documented
<input type="checkbox"/> Medications and Allergies documented	
<input type="checkbox"/> PHQ9 and KID documented	

EVALUATION AND ASSESSMENT	
<input type="checkbox"/> Patient delayed seeing care > 7 days	<input type="checkbox"/> Advanced diagnosis verified
<input type="checkbox"/> Intake information is complete and considered	<input type="checkbox"/> Treatment Plan documented and communicated
<input type="checkbox"/> Chief Complaints identified	<input type="checkbox"/> Informed Consent to provider with patient
<input type="checkbox"/> Physical Exam completed	<input type="checkbox"/> Referral, consultation, and/or communication
<input type="checkbox"/> Diagnostic testing ordered and interpreted	<input type="checkbox"/> Resource Allocation/Case ordered
<input type="checkbox"/> Red flag and contraindications identified	<input type="checkbox"/> Reevaluation and progress
<input type="checkbox"/> Yellow flag identified	

Copyright 2017, Government. All Rights Reserved.

History of Present Illness

Past Family Medical Social History
Review of Systems
Chief Complaints


Copyright 2017, Government. All Rights Reserved.

Physical Comorbidities

Past Medical, Family and Social History

- Prior Major Illnesses and Injuries
- Prior Surgeries
- Prior Hospitalizations
- Current Medications
- Allergies
- Age Appropriate Immunization Status
- Age Appropriate Feeding/Dietary Status
- Marital Status
- Current Employment
- Occupational History
- Alcohol and Tobacco Usage
- Level of Education
- Sexual History


- Ask if there are any members of the patient's family who have had illnesses with features similar to the patient's.
- Determine the health or cause of death of the patient's parents and siblings.
- Establish whether there is a history of heart disease, high blood pressure, cancer, tuberculosis, stroke, diabetes, arthritic conditions, thyroid disease, kidney disease, asthma, blood diseases, sexually transmitted diseases, or any familial diseases.



Copyright © 2021 Elsevier. All Rights Reserved.

Review of Systems

1. Constitutional
2. Eyes
3. Ears, nose, mouth, throat
4. Cardiovascular
5. Respiratory
6. Gastrointestinal
7. Genitourinary
8. Musculoskeletal
9. Integumentary
10. Neurological
11. Psychiatric
12. Endocrine
13. Hematologic/Lymphatic
14. Allergic/Immunologic



Copyright © 2021 Elsevier. All Rights Reserved.


CVA Screening

Has the patient reported any of the following risk factors or symptoms in the medical history?

Is there nausea, vomiting, sensory disturbances (hearing, visual), cramps, weakness, headache, dizziness, and/or loss of consciousness?

Risk Factors:


- Dizziness
- Unsteadiness
- Giddiness
- Vertigo
- Sudden severe pain in the side of the head and/or neck, which is different from any pain the patient has had before
- Age <45 years
- Migraine
- Connective Tissue Disease
- Recent infection (i.e. upper respiratory)



Copyright © 2021 Elsevier. All Rights Reserved.

What are Vital Signs?

These are measurements of the inner workings of the human body and how vital organs, such as the heart and lungs, are functioning.





Copyright © 2021 Elsevier. All Rights Reserved.

INITIAL/PROGRESS VISIT EXAMS


VITAL SIGNS

- HEIGHT
- WEIGHT
- BMI
- BLOOD PRESSURE
- HEART RATE
- RESPIRATION
- BODY TEMPERATURE


Copyright © 2021 Elsevier. All Rights Reserved.

- Height
- Weight
 - Abnormal weight loss or gain
 - Rapid change in height
- BMI (calculated from height/weight)
- Temperature
 - Signs of systemic infection or inflammation in the presence of a fever (temp > 101.4 F or sustained temp > 100.4 F. COVID-19 > 100F).
- Respirations
 - Varies with age, normal reference range is 16-20 breaths/minute.
- Pulse
 - A newborn or infant can have a heart rate of about 130-150 beats per minute.
 - A toddler's heart will beat about 100-120 times per minute.
 - An older child's heartbeat is around 90-110 beats per minute, adolescents around 80-100 beats per minute, and
 - Adults pulse rate is anywhere between 50 and 80 beats per minute.



Copyright © 2021 Elsevier. All Rights Reserved.


- **Normal**
<120 Systolic <80 diastolic
medication not needed, lifestyle
recommendations
- **Pre-hypertensive**
120-139 systolic 80-89 diastolic,
medication not needed, lifestyle
modification (90% chance at 65 to
develop stage 1 and stage 2,
lifestyle changes will decrease risk
to almost 0)
- **Stage 1 hypertension**
140-159 systolic or 90-99
diastolic, lifestyle modifications
given, medications
recommended starting with
thiazide-type diuretics (consider
others if ineffective)
- **Stage 2 hypertension**
>160 systolic or >100 diastolic,
lifestyle modifications given, two-
drug combination therapy
recommended.



Copyright © 2021 Elsevier. All Rights Reserved.

Notes on Blood Pressure


- **Maximum Cuff Pressure** - When the baseline blood pressure is already known or hypertension is not suspected, it is acceptable in adults to inflate the cuff to 200 mmHg and go directly to auscultating the blood pressure. Be aware that there could be an **auscultatory gap** (a silent interval between the true systolic and diastolic pressures).
- **Bell or Diaphragm?** - Even though the Korotkoff sounds are low frequency and should be heard better with the bell, it is often difficult to apply the bell properly in the antecubital fold. For this reason, it is common practice to use the diaphragm when taking blood pressure.
- **Systolic Pressure** - In situations where auscultation is not possible, you can determine systolic blood pressure by palpation alone. Deflate the cuff until you feel the radial or brachial pulse return. The pressure by auscultation would be approximately 10 mmHg higher. Record the pressure indicating it was taken by palpation (60/palp).
- **Diastolic Pressure** - If there is more than 10 mmHg difference between the muffling and the



Copyright © 2021 Elsevier. All Rights Reserved.

Blood Pressure


- Higher blood pressures are normal during exertion or other stress. Systolic blood pressures below 80 may be a sign of serious illness or shock.
- Blood pressure should be taken in both arms on the first encounter. If there is more than 10 mmHg difference between the two arms, use the arm with the higher reading for subsequent measurements.
- It is frequently helpful to retake the blood pressure near the end of the visit. Earlier pressures may be higher due to the "white coat" effect.
- Always recheck "unexpected" blood pressures yourself.



Copyright © 2021 Elsevier. All Rights Reserved.

Pulse, or Heart rate, is the number of times a heart beats per minute (bpm). Heart rates vary by person, and a normal pulse can range between 60 to 100 beats per minute.

Pulse (Heart Rate)





Copyright © 2021 Elsevier. All Rights Reserved.

Pulse

Pulse
Pulse indicates heart rate and it is measured clinically to provide clues to a patient's state of health. It is recorded as beats per minute. Both the rate and the strength of the pulse are important clinically. A high or irregular pulse rate can be caused by physical activity or other temporary factors, but it may also indicate a heart condition.


The pulse strength indicates the strength of ventricular contraction and cardiac output. If the pulse is strong, then systolic pressure is high. If it is weak, systolic pressure has fallen, and medical intervention may be warranted.

Pulse can be palpated manually by placing the tips of the fingers across an artery that runs close to the body surface and pressing lightly. While this procedure is normally performed using the radial artery in the wrist or the common carotid artery in the neck, any superficial artery that can be palpated may be used.





Copyright © 2021 Elsevier. All Rights Reserved.

Pulse



- Note whether the pulse is regular or irregular:
 - **Regular** - evenly spaced beats, may vary slightly with respiration
 - **Regularly Irregular** - regular pattern overall with "skipped" beats
 - **Irregularly Irregular** - chaotic, no real pattern, very difficult to measure rate accurately
- Count the pulse for 15 seconds and multiply by 4.
- Count for a full minute if the pulse is irregular.
- Record the rate and rhythm.



Copyright © 2021 Elsevier. All Rights Reserved.

Staff must report any arrhythmias, irregularities in the pulse rate and pace to the doctor.

Pulse



Copyright © 2021 Elsevier. All Rights Reserved.

Pulse/Blood Pressure in Children

In children, pulse and blood pressure vary with the age. The following table should serve as a rough guide:

Average Pulse and Blood Pressure in Normal Children Age

	Birth	6mo	1yr	2yr	6yr	8yr	10yr
Pulse	140	130	115	110	103	100	95
Systolic	70	90	90	92	95	100	105



Copyright © 2021 Elsevier. All Rights Reserved.

Respiration rate, sometimes referred to as breathing rate, is the number of breaths taken per minute. This measurement is always taken when the individual is at rest.

A single respiration count is equal to the chest rising (inhalation) and falling (exhalation) once. The normal range for an adult is 12 to 28 respirations per minute.

Respiration Rate



Copyright © 2021 Elsevier. All Rights Reserved.

Respiration

<https://youtu.be/nW4sk028uI>

- Best done immediately after taking the patient's pulse. Do **not** announce that you are measuring respirations.
- Without letting go of the patient's wrist begin to observe the patient's breathing. Is it normal or labored?
- Count breaths for 15 seconds and multiply this number by 4 to yield the breaths per minute.
- In adults, normal resting respiratory rate is between 12-28 breaths/minute. Rapid respiration is called tachypnea.



Copyright © 2021 Elsevier. All Rights Reserved.

Temperature is considered normal at 98.6 degrees F (37 degrees C), although anything between 97.6 degrees F (36.4 degrees C) to 99.6 degrees F (37.5 degrees C) is acceptable.

A temperature over 100.4 degrees F (38 degrees C) indicates a fever caused by illness or injury. Hypothermia (low temperature) occurs when the body temperature dips below 95 degrees F (35 degrees C).

Body Temperature



Copyright © 2021 Elsevier. All Rights Reserved.

Temperature

Temperature can be measured in several different ways:

- **Oral** with a glass, paper, or electronic thermometer (normal 98.6F/37C)
- **Axillary** with a glass or electronic thermometer (normal 97.6F/36.3C)
- **Rectal** or "core" with a glass or electronic thermometer (normal 99.6F/37.7C)
- **Aural** (the ear) with an electronic thermometer (normal 99.6F/37.7C)

Of these, axillary is the least and rectal is the most accurate.



Copyright © 2021 Elsevier. All Rights Reserved.

Vital Signs Recap Average Healthy Adults (at rest)

- Blood pressure: 90/60 mm Hg to 120/80 mm Hg
- Respiration: 12 to 18 breaths per minute
- Pulse: 60 to 100 beats per minute
- Temperature: 97.8°F to 99.1°F (36.5°C to 37.3°C)/average 98.6°F (37°C)



Copyright 2021 Elsevier. All Rights Reserved.

Observation

- Observe the patient as they move thru the office, get in and out of the chair, actions while you are performing their history.
- Document what you see:
 - Walks with a limp
 - Difficulty getting out of chair
 - Appears to be in acute pain
 - Medical emergency



Copyright 2021 Elsevier. All Rights Reserved.

Examination

- Observation
 - Gait Analysis
 - Postural Function
- Palpation
- Range of Motion
- Orthopedic Tests
- Neurologic Evaluation
- Vascular Evaluation
- Visceral Evaluation
- X-ray/Lab Evaluation
- External Imaging or Specialty Referral



Copyright 2021 Elsevier. All Rights Reserved.

Be aware of patient's at-risk.
Recognize indications and contraindications for common modalities.
Know Red and Yellow Flags, Contraindications, etc.

At-Risk Patient Population



Copyright 2021 Elsevier. All Rights Reserved.

Red Flags, Yellow Flags, CoMorbidity, and Risk Factors



Copyright 2021 Elsevier. All Rights Reserved.

A serious condition that must be recognized through the history and exam process that typically requires referral to another health care provider

Clinical Red Flags




Copyright 2021 Elsevier. All Rights Reserved.

Red Flags

Immediate Referral

- Fracture/dislocation
 - Significant Trauma
 - Osteoporosis
 - Pathologic Fracture
- Cancer/tumor
 - Night-time Pain
 - Severe Progressive
 - Unexplained Weight Loss
 - Prior History
- Infection
 - Elevated Temperature
 - Night Sweats
 - Intravenous Drug Abuse
 - Immunosuppression
- Vertebrobasilar involvement
- Instability (including degenerative, surgical or rheumatoid etiologies)
- Progressive scoliosis
- Severe osteoporosis
- Severe hypertension
- Vertebrobasilar involvement
- Visceral pathology
- Inflammatory Arthritides
- Cauda Equina Syndrome (loss of bladder/bowel function)



Copyright © 2021 Osseointegration. All Rights Reserved.

Initial Visit Patient Safety Checklist

Use this checklist to gather essential information regarding the initial or re-admission of a patient to determine if the patient is at risk for an adverse event and there is a need for the monitoring of key patient safety activities.

PATIENT INTAKE AND INFORMATION GATHERING


<input type="checkbox"/> Intake forms completed and signed	<input type="checkbox"/> Patient is showing unsteadiness in movements
<input type="checkbox"/> Patient's mental status is not normal	<input type="checkbox"/> Patient has fallen within the last 6 months
<input type="checkbox"/> Patient recently had a hospitalization or surgery	<input type="checkbox"/> Patient is on medication or has allergies

PRE-EVAL EHR DOCUMENTATION

<input type="checkbox"/> Demographics completed	<input type="checkbox"/> Visit signs documented and communicated
<input type="checkbox"/> Referring Physician identified and documented	<input type="checkbox"/> Potential Clinical Red Flags communicated
<input type="checkbox"/> Medications and Allergies documented	
<input type="checkbox"/> PHQ-9 and RSI documented	

EVALUATION AND ASSESSMENT

<input type="checkbox"/> Patient delayed seeking care (> 7 days)	<input type="checkbox"/> Achieved diagnosis certainty
<input type="checkbox"/> Intake information is complete and condensed	<input type="checkbox"/> Treatment Plan documented and communicated
<input type="checkbox"/> Chief Complaints identified	<input type="checkbox"/> Informed Consent by provider with patient
<input type="checkbox"/> Physical Exam completed	<input type="checkbox"/> Referral, collaboration, and/or communication
<input type="checkbox"/> Diagnostic testing ordered and interpreted	<input type="checkbox"/> Outcome Assessment Tools selected
<input type="checkbox"/> Red flags and contraindications reviewed	<input type="checkbox"/> Re-evaluation and progress
<input type="checkbox"/> Yellow flags identified	




Copyright © 2021 Osseointegration. All Rights Reserved.

Treatment




Copyright © 2021 Osseointegration. All Rights Reserved.

Evidence-Informed Practice




The Evidence-based Medicine Triad
 Source: Florida State University College of Medicine



Copyright © 2021 Osseointegration. All Rights Reserved.

Standard of Care

How does your state licensing board view YOUR responsibilities as a clinician, within the interest of public safety?



Copyright © 2021 Osseointegration. All Rights Reserved.

Clinical Competencies

Efficacious Treatment Approaches
 Competency of Doctor and Staff in delivery of services



Copyright © 2021 Osseointegration. All Rights Reserved.

Are you and your staff attending regular clinical education training?
 Do you provide hands-on training for staff?
 Are you using FDA approved devices?
 Does your treatment follow guidelines?
 Are you monitoring and documenting the progress of your patients?

Questions to Ask



Copyright © 2021 ChiroSource. All Rights Reserved.

Treatment Visit Screening

Treatment Visit Patient Safety Checklist
 Use this checklist to gather essential information regarding the Treatment Visit of a patient to determine if the patient is at risk for an adverse event and there is a need for the monitoring of long patient safety studies.

TREATMENT VISITS	
<input type="checkbox"/> Patient is showing unsteadiness in movements	<input type="checkbox"/> Document subjective and objective findings
<input type="checkbox"/> Patient recently had a hospitalization or surgery	<input type="checkbox"/> Assess procedure and patient response
<input type="checkbox"/> Change in vital signs measured/communicated	<input type="checkbox"/> Assess progress of the patient
<input type="checkbox"/> Significant event identified requiring evaluation	<input type="checkbox"/> Assess diagnostic certainty
<input type="checkbox"/> Patient has a change in medication or allergy	<input type="checkbox"/> Patient monitoring after treatment
<input type="checkbox"/> Patient has fallen within the last 6 months	<input type="checkbox"/> Potential Clinical Red Flags communicated
<input type="checkbox"/> Patient's mental status is not normal	

POST-EVAL/TREATMENT EHR DOCUMENTATION	
<input type="checkbox"/> Patient findings/visit information documented	<input type="checkbox"/> Patient response monitored and reported
<input type="checkbox"/> Patient progress assessed and documented	<input type="checkbox"/> Therapeutic activities/modifications prescribed
<input type="checkbox"/> Procedures performed correctly without incident	<input type="checkbox"/> Coding and charges completed as followed



Copyright © 2021 ChiroSource. All Rights Reserved.

Misinformed Treatment Plans

Communicating to patients regarding the treatment plan and expectations of care process.



Copyright © 2021 ChiroSource. All Rights Reserved.

Care Management Considerations

Transitional Care (Hand-off)
 Environment/Falls
 Medication Errors/Reconciliation
 Team/Communication




Copyright © 2021 ChiroSource. All Rights Reserved.

Dry Needling/Acupuncture Adverse Effects

The act of puncturing the skin comes with a number of predictable adverse events (bruising or bleeding, pain during or following treatment) which commonly occur and are mild in nature.


This may be considered normal side effects of treatment. However, from the patient's perspective they may be considered adverse particularly if the patient has not been educated about the risks associated with their dry needling/acupuncture technique.



Copyright © 2021 ChiroSource. All Rights Reserved.

Manipulation/Manual Therapy Potential Risks

- ✓ Temporary soreness or increased symptoms or pain It is not uncommon for patients to experience temporary soreness or increased symptoms or pain after the first few treatments.
- ✓ Dizziness, nausea, flushing These symptoms are relatively rare. It is important to notify the doctor if you experience these symptoms during or after your care.
- ✓ Fractures When patients have underlying conditions that weaken bones, like osteoporosis, they may be susceptible to fracture. It is important to notify your doctor if you have been diagnosed with a bone weakening disease or condition. If your doctor detects any such condition while you are under care, you will be informed, and your treatment plan will be modified to minimize risk of fracture.
- ✓ Disc herniation or prolapse Spinal disc conditions like bulges or herniations may worsen even with chiropractic care. It is important to notify your doctor if symptoms change or worsen.
- ✓ Stroke According to the most recent research, there is no evidence of excess risk of stroke associated with chiropractic care. Regarding neck pain and headache symptoms, there is an association between stroke and visits to all provider-types, including primary care medical visits, which may occur before or during the provider visit.
- ✓ Other risks associated with chiropractic treatment include rare burns from physiotherapy devices that produce heat.
- ✓ Bruising Instrument assisted soft tissue manipulation may result in temporary soreness or bruising.



Copyright © 2021 ChiroSource. All Rights Reserved.

Chiropractic Clinical Assistant Competency

- Formal training completion with testing
- Understand supervision rules for your state
- Patient response
- Doctor communication - orders



Copyright 2021 ChiroSource. All Rights Reserved.

Recognizing and Preventing Safety Hazards

1. Therapy Modalities
2. Hydraulic/Spring-loaded adjusting tables
3. Sharps (i.e. needles) and Sharps Containers
4. Theraband/Exercise Stations



Copyright 2021 ChiroSource. All Rights Reserved.

Therapeutic Modalities and Table Equipment

- Are all therapeutic modalities and equipment (both, company and employee-owned) used by staff, providers and workforce members at their workplace in good condition?
- Are all of the operating manuals and instructions available to staff, providers and workforce members for all therapeutic modalities and equipment?
- Are staff, providers and workforce members made aware of the hazards caused by faulty or improperly used modalities and equipment?
- Are all cord-connected, electrically operated modalities and equipment effectively grounded or of the approved double insulated type?
- Are children monitored at all times and parent/guardian warned of crush risk or safety issue around modalities?



Copyright 2021 ChiroSource. All Rights Reserved.

Therapeutic Modalities and Table Equipment

- Are all therapeutic modalities and equipment turned off after use and remain off prior to patient use?
- Do patients know what to expect prior to the application of the modality?
- Do patients know what to expect as potential temporary symptoms or reactions to the application of the therapy?



Copyright 2021 ChiroSource. All Rights Reserved.

Theraband Exercise Station

Eye Protection



Copyright 2021 ChiroSource. All Rights Reserved.

Re-Cap



Copyright 2021 ChiroSource. All Rights Reserved.

Clinical Conscientiousness and Awareness

Maintaining your clinical mindset



Copyright © 2021 Cleveland. All Rights Reserved.

The "Walk"

Welcome
Ask
Listen
Knowledge



Copyright © 2021 Cleveland. All Rights Reserved.

Screening Patients:

Why are you here today?
Has there been a change in how you are feeling since your last visit?
Have you seen anyone else about your health?
Do you have questions about...
Are you worried about your health?

Situational Awareness:

No change or worsening
Observation of patient's mental status, behaviors, or characteristics

Has there been a "Significant Event"?



Copyright © 2021 Cleveland. All Rights Reserved.

Patient Safety Questionnaire

Patient Safety Questionnaire

Please answer the following questions as it pertains to your visit today - we will use this information as we strive to help you!

- | Yes | No | |
|--------------------------|--------------------------|---|
| <input type="checkbox"/> | <input type="checkbox"/> | I have felt unsteady and off balance when I walk, stand or move. |
| <input type="checkbox"/> | <input type="checkbox"/> | I have recently had a hospitalization or surgery since my last visit. |
| <input type="checkbox"/> | <input type="checkbox"/> | There has been a change in my condition since my last visit. |
| <input type="checkbox"/> | <input type="checkbox"/> | Something happened to me since my last visit (i.e., accident, fall, etc.) |
| <input type="checkbox"/> | <input type="checkbox"/> | I have had a recent change in my medication or have a new allergy. |
| <input type="checkbox"/> | <input type="checkbox"/> | I have fallen within the last 6 months. |
| <input type="checkbox"/> | <input type="checkbox"/> | My mental processing or thinking doesn't seem very clear or is different to me. |
| <input type="checkbox"/> | <input type="checkbox"/> | I need to talk to the doctor about my progress. |
| <input type="checkbox"/> | <input type="checkbox"/> | I am concerned about the treatment I am getting. |
| <input type="checkbox"/> | <input type="checkbox"/> | I have questions I need to ask the doctor today. |



Copyright © 2021 Cleveland. All Rights Reserved.

Does the patient's clinical
presentation require urgent need
for evaluation and/or care?

The doctor must be informed of any new information
about the patient that has been related to staff.



Copyright © 2021 Cleveland. All Rights Reserved.



Stay Connected to Established Patients
who are under a treatment plan.

Following the treatment plan, evidence-informed
care guidelines, and the patient's response to care...



Copyright © 2021 Cleveland. All Rights Reserved.

It just takes one thing to block the incident...



<https://www.youtube.com/watch?v=7791e6Cv8K4>

Copyright © 2011, Clark Howard. All Rights Reserved.



Thank you!

Scott Munsterman, DC, FICC, CPCO



Copyright © 2011, Clark Howard. All Rights Reserved.